

Effects of Berberine on Leukemia with a Focus on Its Molecular Targets

Mohammad Reza Mazandarani¹, Parisa Maleki Dana², Zatollah Asemi², Jamal Hallajzadeh^{3,*}, Mohammad Ali Mansournia⁴ and Bahman Yousefi^{5,6}

¹Department of Community Nutrition, Faculty of Nutrition and Food Technology, Shahid Beheshti University of Medical Sciences and Health Services, Tehran, Iran; ²Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, I.R. Iran; ³Department of Biochemistry and Nutrition, Research Center for Evidence-Based Health Management, Maragheh University of Medical Sciences, Maragheh, Iran; ⁴Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran; ⁵Molecular Medicine Research Center, Tabriz University of Medical Sciences, Tabriz, Iran; ⁶Department of Biochemistry, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

ARTICLE HISTORY

Received: March 18, 2021
Revised: October 04, 2021
Accepted: December 01, 2021

DOI:
10.2174/1871520622666220324092302



CrossMark

Abstract: Leukemia is common among both women and men worldwide. Besides the fact that finding new treatment methods may enhance the life quality of patients, there are several problems that we face today in treating leukemia patients, such as drugs' side effects and acquired resistance to chemotherapeutic drugs. Berberine is a bioactive alkaloid found in herbal plants (e.g., Rhizoma coptidis and Cortex phellodendri) and exerts several beneficial functions, including anti-tumor activities. Furthermore, berberine exerts antiproliferative and anti-inflammatory effects. Up to now, some studies have investigated the roles of berberine in different types of leukemia, including acute myeloid leukemia and chronic lymphocytic leukemia. In this review, a detailed description of the roles of berberine in leukemia is provided. We discuss how berberine involves different molecular targets (e.g., interleukins and cyclins) and signaling pathways (e.g., mTOR and PI3K) to exert its anti-tumor functions and how berberine is effective in leukemia treatment when combined with other therapeutic drugs.

Keywords: Berberine, leukemia, apoptosis, proliferation, autophagy, inflammation.

1. INTRODUCTION

Based on the American Cancer Society statistics, the estimated new cases of leukemia and its estimated deaths in 2021 are reported to be 61,090 and 23,660 respectively [1]. There are different types of leukemia, but four of them are more common, including myelodysplastic syndrome, myeloproliferative neoplasm, chronic myeloid leukemia (CML), and acute myeloid leukemia (AML) [2, 3]. CML, a malignant clonal hematological disease, is defined by the myeloid neoplastic cells spread into the blood circulation and bone marrow [4]. Another common type of leukemia is chronic lymphocytic leukemia (CLL), which is the disease of the elderly with a median age of 72 years [5]. Leukemia risk factors are classified into four categories: familial, genetic, environmental, and lifestyle risk factors. Benzene, formaldehyde, 1,3 butadiene, high dose ionizing radiation, chemotherapeutic drugs, and electromagnetic fields are some of the environmental risk factors. Smoking, obesity, and dietary intake are some of the lifestyle risk factors [6, 7]. 5-year relative survival of patients varies from type to type. While 5-year survival is 27.4% in AML patients, this number rises to 84.2% for CLL patients [8]. Besides, more than half of patients surviving the disease relapse after treatment [9]. Therefore, new treatment modalities are essential for patients with leukemia.

Berberine, a phytochemical compound, is mostly extracted from roots, rhizome, and stem bark of barberry [10]. In recent years, berberine has been discovered to have a variety of pharmacological and biological activities such as antimicrobial, antihelminthic, anti-inflammatory, and anti-oxidative effects [11].

Furthermore, several studies have investigated the anti-cancer activities of berberine [11]. In this review, a detailed description of berberine roles in leukemia is provided. We discuss how berberine involves different molecular targets and signaling pathways to exert its anti-tumor functions.

2. PATHOGENESIS OF LEUKEMIA

In acute leukemia, the abnormal blood cells are usually blasts that remain immature. Consequently, fail to perform their expected functions. Acute leukemia exhibits fast development and requires immediate attention. In chronic leukemia, there are also some blast cells. However, these blast cells are different from cells presented in acute leukemia. Indeed, they are more mature and may still have their normal functions [12]. The progression of chronic leukemia is usually much slower than acute leukemia and may not need immediate treatment. Based on another type of classification, leukemia can be divided into two types according to the affected cells, lymphoblastic leukemia and myeloid leukemia. In lymphoblastic leukemia, the malignant changes involve a type of bone marrow cells that eventually become lymphocytes. In myeloid leukemia, the malignant changes involve the type of bone marrow cells that eventually become red blood cells, some other types of white cells, and platelets [12]. These two types of divisions lead to four major subtypes in total, including ALL, CLL, AML, and CML. CML is a stem cell disorder characterized by the proliferation of hematopoietic stem cells due to the BCR-ABL fusion oncogene, resulting in reciprocal translocation t(9;22) (q34;q11). It is known that imatinib, a tyrosine kinase inhibitor that blocks the breakpoint cluster region protein/ Abelson murine leukemia viral oncogene homolog (BCR-ABL1) leads to the suppression of PI3K/AKT, JAK/STAT, and RAS/RAF/MEK/ERK pathways [3]. CLL is an incurable neoplastic disorder that is the most common form of leukemia. It is character-

*Address correspondence to this author at the Department of Biochemistry and Nutrition, Research Center for Evidence-Based Health Management, Maragheh University of Medical Sciences, Maragheh, Iran;
E-mail: jamal.hallaj@yahoo.com

ized by a gradual accumulation of small and mature B cells with typical B-cell markers, including CD5, CD19, CD23, and CD20 [13]. ALL is caused by the accumulation and proliferation of lymphoid progenitor cells in different tissues, including bone marrow [14]. In AML, myeloid stem cells are abnormally proliferated and differentiated. What changes the normal maturation of myeloid precursors can be some translocations in the chromosome, including t(8:21) in core-binding factor AML [15]. Treatment for AML is chemotherapy with cytarabine and anthracycline or hypomethylating 0.agents and allogeneic hematopoietic stem cell transplant [16, 17].

3. OXIDATIVE STRESS IN LEUKEMIA

Oxidative stress results from a biochemical imbalance consisting of excessive production of reactive oxygen species (ROS) and nitrogen species, leading to oxidative damage to biomolecules and cannot be defused by anti-oxidative systems [18]. ROS are a group of heterogeneous molecules that are generated by mature myeloid cells throughout innate immune responses. The imbalance in production and degradation of ROS is the definition of oxidative stress which has been observed in some malignancies such as leukemia [19, 20]. The activity of ROS is a major part of cell function. While irregularities in ROS levels result in DNA impairment. Moreover, higher levels of ROS cause changes in the metabolic state of cells [21]. ROS role in oxidative stress is through influencing intracellular cascades such as mitogen-activated protein kinases (MAPKs), phosphatidyl inositol-3-kinase (PI3K/Akt), phospholipase C-g1, protein kinase C, nuclear factor- κ B (NF- κ B), and Janus kinase/signal transducer and activator of transcription (JAK/STAT). ROS prompts the expression of heat shock proteins (Hsp27), immediate early genes (c-jun and c-fos), and hypoxia-inducible factors. Also, ROS affects antioxidant enzymes, which are involved in redox homeostasis, transforming oncoproteins, and growth factors [22].

4. INFLAMMATION AND LEUKEMIA

Inflammation and inflammatory cytokines have always been a major reason for the onset and progression of leukemia [23]. Pro-Inflammatory agents like IL-1, IL-6, IL-15, IL-17, IL-23, and TNF α have a role in leukemia severity. On the other hand, anti-inflammatory mediators like IL-4, IL-10, IL-13, transforming

growth factor (TGF β), and interferon α (IFN α) seem to inhibit the progression of leukemia. These complex interactions among pro- and anti-inflammatory cytokines make a carcinogenic microenvironment that influences cell proliferation and survival in leukemia [13, 24]. The major function of TNF α is to activate NF- κ B. Meanwhile, IL-10 deactivates NF- κ B, which promotes pro-inflammatory cytokine TNF α , IL-6, and IL-12 [13]. IL-6 can trigger the expression of miR-21, which is an inflammatory stimulus. TNF α and IFN- β stimulate the expression of miR-155 and its increased levels are found in the bone marrow of leukemia patients and cause hyperproliferation of B-cells [13].

5. BERBERINE

Berberine (chemical formula: C₂₀H₁₈NO₄) is a bioactive isoquinoline alkaloid that exists in herb plants, such as Cortex phellodendri and Rhizoma coptidis [25, 26]. Rutaceae, Berberidaceae, and Papaveraceae are other sources of berberine. Berberine was at first used as a broad-spectrum antibacterial drug. Pharmacological uses such as antibacterial, anti-inflammatory, antihypertensive, hypolipidemic, and antidiarrheal effects were the main applications of berberine in the past [27]. Berberine is also an antioxidant and anti-inflammatory agent. By exerting anti-oxidative and anti-inflammatory functions, berberine could be effective in type 2 diabetes and cardiovascular diseases' management (Fig. 1) [28]. Berberine affects the development of tumor cells by inhibiting the growth of tumor cells as well as stimulating apoptosis and cell cycle arrest [29-33]. Berberine lowers plasma LDL by increasing LDLR mRNA expression at a post-transcriptional level [34]. The main organ for berberine metabolism is the liver. The major cytochrome P450 (CYPs) for producing berberine metabolites is CYP2D6. CYP1A2, 3A4, 2E1, and CYP2C19 are enzymes that are also involved in berberine metabolism [35, 36]. It has been discovered that berberine could increase the activity of AMP-activated protein kinase, which leads to phosphorylation of the tumor suppressor gene p53 in vascular smooth muscle cells as well as growth inhibition and apoptosis in non-small cell human lung cancer cells [11]. Besides the advantages of berberine, there are some limitations, such as poor aqueous solubility, slight absorption, and low bioavailability. To overcome these limitations, the utilization of nanotechnology-based platforms (such as nanoparticles, hydrogels, and micelles) and various analogs are suggested (Fig. 1) [32].




Fig. (1). Different advantages and disadvantages of berberine. (A higher resolution / colour version of this figure is available in the electronic copy of the article).

6. ANTI-INFLAMMATORY AND ANTIOXIDANT EFFECTS OF BERBERINE

Several studies have investigated the anti-inflammatory role of berberine in various diseases. Berberine is reported to reduce the expression levels of various pro-inflammatory cytokines, including IL-6, IL-8, IL-13, IFN- γ , and TNF- α [37]. Berberine can also modulate the tight junction damage of intestinal epithelium, which is induced by pro-inflammatory cytokines [38, 39]. In chronic respiratory diseases, berberine exerts anti-inflammatory effects *via* different signaling pathways, including p38 MAPK, NF- κ B, and ERK1/2 [40]. Moreover, in non-alcoholic fatty liver disease of rats, berberine is reported to suppress the inflammatory response by regulating the Angptl2 pathway [41]. Therefore, berberine attenuates inflammatory processes through various mechanisms.

Berberine is shown to increase the hepatic antioxidant enzymes in thioacetamide-induced liver fibrosis, leading to the suppression of hepatic fibrosis [42]. Another study has reported that berberine exerts protective effects on PC-12 cells against oxidative damage by suppressing ROS through PI3K/AKT/mTOR pathway [43]. Berberine is capable of enhancing the activities of glutathione peroxidase and catalase enzymes in mice intestinal tissue, resulting in reduced oxidant status [44]. Moreover, berberine modulates the formation of ROS in spiral ganglion cells [45]. Altogether, these pieces of information indicate that berberine is a potential antioxidant agent which enhances antioxidant status by different mechanisms [46].

7. BERBERINE AND LEUKEMIA

Findings have shown that berberine exerts multiple functions in leukemia; thus, it can be used as an antitumor agent for treating this cancer. However, the exact underlying mechanisms of berberine's function in leukemia are to be elucidated. Herein, we take a look at the details provided by research until now, aiming to provide an insight into which areas are not paid attention to enough and which areas are well-understood.

Development and progression of tumors as well as their malignant features, such as changes in differentiation of cells, instabilities in the genome, and increased metastatic abilities, have been related to P53 dysfunction [47, 48]. Mutation of TP53 happens in approximately 8% of de novo AML cases. Besides, Li-Fraumeni syndrome-related AML is also rare. Thus, other p53 abnormalities that are non-mutational may be involved in AML and other types of leukemia. In fact, evidence has shown that non-mutational dysfunction of wtp53 may happen in almost all subsets of AML [49]. Mouse double minute-2 (MDM2) is a regulator of P53, which provides stability and cellular localization of P53. The interaction between P53 and MDM2 induces proteasomal degradation of P53, leading to an inhibition of its transcriptional activity [50, 51]. Some studies indicate that MDM2 can be a potential target for treating ALL [52]. In a study on P53-null leukemic cells, berberine has been shown to reduce the mRNA level of MDM2 as well as increasing self-ubiquitination of MDM2 [53]. While 100 μ M of berberine induces autophagy and apoptosis in leukemic cells, 3-methyladenine, which is an inhibitor of autophagy, reverses the berberine effect on autophagy [53]. Besides, forcing cells to over-express MDM2 has an unfavorable impact on berberine-induced effects, such as autophagy, apoptosis, and doxorubicin-related effects [53]. Noteworthy, it is suggested that MDM2 reverse effects on berberine are not dependent on P53 [53]. Suppression of XIAP by berberine in p53-null leukemia cells leads to downregulation of XIAP protein and inhibition of MDM2 expression [54]. Moreover, western blot analysis showed that XIAP suppression by berberine enhances doxorubicin-induced apoptosis [54]. On the other hand, another study has shown that berberine apoptotic effects are related

to MDM2 and P53 [55]. Contrary to doxorubicin, which results in the activation of P53 and MDM2 upregulation, berberine downregulates MDM2 besides the activation of P53 [55]. Moreover, berberine-induced downregulation of MDM2 has been linked to the death domain associated protein (DAXX) [55]. It is also suggested that berberine can be beneficial in the chemo-resistant form of leukemia since these cells overexpress MDM2 [55].

Proviral integration of Moloney virus-2 (PIM-2) is an anti-apoptotic factor and a member of the serine/threonine kinase family [56]. Cytokines, which are involved in hematopoietic cells maturation, are responsible for the regulation of PIM-2 expression [57]. Evidence reveals that PIM-2 is overexpressed in AML and ALL [56]. Overexpression of PIM-2 has been associated with overall survival in patients with AML [56]. Furthermore, PIM-2 downregulation inhibits the proliferation of leukemia cells [58]. PIM-2 is also a direct target of miR-24-3p [59]. In a recent study, Wang *et al.* [59] reported that treating cells with 50 μ M of berberine for 24 hours can affect PIM-2 by increasing miR-24-3p. It is indicated that berberine plays anti-tumor roles by inducing apoptosis and reducing the viability of ALL cells through downregulating XIAP in P53-null and P53-mutant cells, as shown by qRT-PCR. Besides, berberine effects on apoptosis are reversed by blocking miR-24-3p or PIM-2 [59].

As we mentioned earlier, imatinib is a tyrosine kinase inhibitor that is used for the treatment of CML by reducing BCR-ABL transcripts [60]. However, it is observed that patients may develop resistance to imatinib during their treatment course due to the sub-clones which bear BCR-ABL mutations [61]. Berberine is shown to bind to a domain of BCR-ABL, which is called protein tyrosine kinase (PTK) [62]. Consequently, both normal BCR-ABL transcripts and BCR-ABL transcripts with the mutation in T315I are degraded through the autophagic lysosome pathway [62]. Additionally, LRSAM1, an E3 ubiquitin-protein ligase, is involved in the mentioned process [62]. Therefore, berberine could be an effective antitumor agent in CML patients who are sensitive or resistant to imatinib. Studies implied that targeting Aryl Hydrocarbon Receptor (AHR) signaling can be useful in treating AML since this pathway is suppressed in leukemic stem cells [63]. AHR, a ubiquitous helix-loop-helix transcription factor, plays various roles in biological processes such as inflammation and cell division [64]. Furthermore, it is associated with the proliferation of normal hematopoietic progenitors. UT-7 cells that express BCR-ABL have been shown to express lower levels of AHR [64]. Berberine which is considered an Ahr ligand, results in the overexpression of AHR target genes (CYP1A1 and IL1 β), as evidenced by RT-PCR. Consequently, this leads to inhibition of monocytes proliferation [65]. Berberine also increases IL-10, downregulates Cdk2, and upregulates p21, p27, and p53 genes in the AML cell line [65].

Knockdown of nucleophosmin/B23 is shown to downregulate Akt/mTOR signaling pathway, leading to modulation of drug resistance [66]. This ubiquitous nuclear phosphoprotein plays various roles in cells, such as progression of the cell cycle, ribosome biogenesis, duplication of the centrosome, and cell growth [67]. Overexpression of nucleophosmin/B23 leads to the proliferation of tumor cells, differentiation blockade, and resistance to apoptosis [67]. It is suggested that nucleophosmin/B23 may have a role in controlling the cellular response to apoptosis induction in human leukemia HL-60 cells [68]. Berberine can induce apoptosis by downregulating the nucleophosmin/B23 and telomerase activity [68]. Studies conducted on rats report that berberine inhibits the release of β -hexosaminidase, histamine, IL-4, and TNF- α in basophilic leukemia cells (RBL-2H3 cells) [69]. Berberine induces arrest of the cell cycle at the G1 phase and G2/M phase in leukemia, like in other cancers [25, 26, 70]. The cell cycle arrest at the G2/M phase is sim-

ultaneous with increased levels of Wee1 and 14-3-3 σ and also decreased levels of Cdc25c, CDK1, and cyclin B1 [70]. Berberine has anti-proliferative actions in leukemia cells that are implemented by inducing chromatin condensation, DNA fragmentation and triggering the activation of PARP, caspase-3, and caspase-8 without activating caspase-9 [71-73]. Berberine also inhibits the proliferation of Jurkat cells. It causes cell cycle arrest at the G1 phase *via* downregulation of cyclin D1, cyclin E, and CDC2. Berberine induces apoptosis by regulating the ephrin-B2 and VEGFR2 signaling, along with modulating the MEK/ERK and PTEN/PI3K/AKT/mTOR signaling simultaneously [65]. Berberine's anti-inflammatory effects are through MAPK and COX-2 signaling pathways, where it reduces the expression of COX-2, which leads to apoptosis in T-cell ALL. Berberine suppresses IL-2 mRNA expression and protein secretion [25, 72]. By using transwell migration chambers, it is shown that berberine has a prohibitory role in tumor cell migration in a dose-dependent manner. SDF-1, a homeostatic chemokine that signals through CXCR4, is expressed by hematopoietic tumor cells. Berberine partly inhibits AML cells migration by reducing SDF-1 protein [74]. The apoptotic effects of berberine can be done by increasing ROS and Ca²⁺ production. Furthermore, berberine decreases mitochondrial membrane potential, which leads to the release of cytochrome c and the cleavage of procaspase-3 [75]. Studies conducted *in vitro* and *in vivo* revealed that berberine inhibits the expression of NAT1 protein by affecting the mRNA in HL-60 cells [76-78].

8. STRUCTURE-RELATED ANTITUMOR ACTIVITIES OF BERBERINE

Despite berberine's various beneficial activities, there are some obstacles to applying it as an antitumor agent, such as its difficulty in penetrating cytomembrane and low lipid solubility [79-83]. Poor bioavailability is another drawback in berberine application [82]. Thus, some changes have been made to berberine in investigations for modifying and transforming its chemical structure. C-9 and C-13 are the main locations where structural changes are made to

enhance the anti-cancer effects of berberine [84-91]. Accordingly, several studies have been conducted on berberine derivatives to assess the associations between structural modifications and anti-tumor functions [46].

Regarding leukemia, Okubo *et al.* [73] showed that a methylenedioxy group on ring A is essential for better anti-cancer activities of berberine compounds since the compounds lacking this group were weaker in inhibitory activities. Furthermore, the alkylation of the hydroxyl group at C-9 position makes berberine compounds more effective.

CONCLUSION

Leukemia is a common type of cancer worldwide. Besides the fact that finding new treatment methods may enhance the life quality of patients, there are several problems that we face today in treating leukemia patients, such as drugs' side effects and acquired resistance to chemotherapy drugs. Evidence shows that berberine can be used to treat different diseases, including cancer, due to its ability to involve different signaling pathways. It is demonstrated that berberine has anti-tumor effects on various cell lines, such as induction of apoptosis, inhibiting proliferation, suppressing metastasis, anti-inflammatory activities, and induction of autophagy.

Studies suggest that berberine is a potentially effective agent for inhibiting the development and progression of leukemia (Table 1 and Fig. 2). Berberine exerts this inhibitory role by involving several molecules and signaling pathways. Induction of apoptosis and autophagy, as well as inhibiting proliferation, are the main mechanisms by which berberine is considered a beneficial target for managing leukemia. Noteworthy, studies have shown that treating cells with berberine is not limited to one type of leukemia and it affects different leukemia cell lines, including Jurkat cells, EU-4, and HL-60. These functions of berberine on leukemia cell lines seem to involve several mechanisms that require further exploration. Studies concerning the berberine anti-tumor effects have some limitations. First of all, the underlying anticancer roles are not fully understood. Secondly, most of the studies are *in vitro*; thus, *in vivo*




Fig. (2). Schematic representation of how berberine exerts its functions in leukemic cells. (A higher resolution / colour version of this figure is available in the electronic copy of the article).

Table 1. Studies investigated the roles of berberine in different types of leukemia.

<i>In vivo/ In vitro</i>	Dosage	Cell Line(s)	Findings	Refs.
<i>In vitro</i>	20 µg/ml	Jurkat	Berberine played its anti-inflammatory role possibly due to suppressing p38 MAPK expression	[25]
<i>In vitro</i>	6-100 µM	HL-60	Berberine iodide and acetoneberberine presented apoptosis-inducing activity and tumor-specific cytotoxicity	[92]
<i>In vitro</i>	59.2 µg/ml	Jurkat	Berberine induced apoptosis and S phase arrest of cell cycle	[93]
<i>In vitro</i>	NA	K562	Berberine resulted in apoptosis	[94]
<i>In vitro</i>	75 µM/ml	K562	It is suggested that berberine may play its apoptosis-inducing role by reducing survivin levels	[95]
<i>In vitro, In vivo</i>	100 µM	Jurkat, U937	Berberine induced apoptosis and autophagy as well as promoting doxorubicin-induced autophagy	[53]
<i>In vitro</i>	100 µM	EU-4 cells	Berberine led to apoptosis which is mediated by XIAP suppression	[54]
<i>In vitro</i>	100 µM	EU-1, EU-4, EU-6, EU-8, Sup-B13, UOC-B1	Berberine induced apoptosis mediated by P53 and MDM2	[55]
<i>In vitro, In vivo</i>	50 µM	KOPN-8, EU-4, NALM-6, EU-6, SEM	Berberine induced apoptosis and reduces cell viability through XIAP downregulation	[59]
<i>In vitro, In vivo</i>	5 µM	-	Berberine stimulated autophagic degradation of BCR-ABL and BCR-ABL T315I and overcomes imatinib-resistance	[62]
<i>In vitro</i>	50 µg/ml	HL60	Berberine induced apoptosis which is related to downregulation of nucleophosmin/B23 and telomerase activity	[68]
<i>In vitro</i>	15-60 µM	HL-60, murine WEHI-3	Berberine induced G2/M arrest by inhibiting cyclin B1 and promoting Wee1	[70]
<i>In vitro</i>	50 µM	HL-60	Berberine activated caspase-3 and caspase-8 leading to apoptosis	[73]
<i>In vitro</i>	60 µM	HL-60, murine WEHI-3	Berberine induced apoptosis through activating caspase-3	[75]
<i>In vitro</i>	40-160 µM	L1210	Berberine inhibited the NAT activity and AF-DNA adduct formation	[77]
<i>In vitro</i>	1.0, 2.5, 4.0 and 5.0 mM	HL60	Berberine-derived Quinolono-benzo-[5, 6]-dihydroisoquinolium compounds are highly selective ligands for G-quadruplex DNA in c-myc oncogene	[96]
<i>In vitro</i>	25-100 µg/ml	L1210	Berberine led to G0/G1 cell cycle arrest and apoptosis	[97]
<i>In vitro</i>	6.25 µM	Jurkat	Berberine had a synergistic effect with TPD7 on cell growth through Ephrin-B2 signaling	[65]
<i>In vivo</i>	200mg/Kg	WEHI-3	Berberine inhibited leukemic cell line	[98]
<i>In vitro</i>	25 µg/ml	HL-60	Berberine induced apoptosis	[99]
<i>In vitro</i>	4 microg/ml	L120	Berberine induced apoptosis	[100]
<i>In vitro</i>	15, 57, 74, 221, 40 and 80 µM/mL (IC50)	K562, U937, P3H, Raji	Berberine and coptisine which are compounds of Coptis chinensis inhibited cell proliferation	[1]
<i>In vitro</i>	100 µM	THP-1	Berberine increased the expression of CDK inhibitors and anti-inflammatory cytokines as well as downregulating CDK2.	[65]
<i>In vitro, In vivo</i>	100 µM	EU-6 and SKW-3	Berberine induces autophagic death by inactivating AKT/mTORC1 pathway	[101]
<i>In vitro, In vivo</i>	10 µmol/L	K562	Berberine induces autophagic degradation of BCR-ABL T315I and BCR-ABL by recruiting LRSAM1 to overcome imatinib resistance	[102]

and clinical investigations must be conducted to confirm the observed effects *in vitro*.

LIST OF ABBREVIATIONS

ALL = Acute Lymphoblastic Leukemia
 CLL = Chronic Lymphoblastic Leukemia
 AML = Acute Myeloid Leukemia
 CML = Chronic Myeloid Leukemia
 ROS = Reactive Oxygen Species
 MAPK = Mitogen-activated Protein Kinase

PI3K = Phosphatidyl Inositol-3-Kinase
 NF-κB = Nuclear Factor-κB
 JAK/STAT = Janus Kinase/signal Transducer and Activator of Transcription
 COX-2 = Cyclooxygenase-2
 Hsp = Heat Shock Proteins
 AhR = Aryl Hydrocarbon Receptor
 MDM2 = Mouse Double Minute-2
 XIAP = X-linked Inhibitor of Apoptosis Protein
 PIM-2 = Proviral Integration of Moloney Virus-2

DAXX	=	Death Domain Associated Protein
BCR-ABL	=	Breakpoint Cluster Region Protein/Abelson Murine Leukemia Viral Oncogene Homolog
AHR	=	Aryl Hydrocarbon Receptor

AUTHORS' CONTRIBUTIONS

JH and MRMF contributed to the conception, design, and drafting of the manuscript. ZA, PMD, MAM, and BY contributed to reviewing relevant literature. All authors approved the final version for submission.

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors would like to thank Shohada Clinical Research Development Unit, Shohada Hospital, Tabriz University of Medical Science, Tabriz, Iran.

REFERENCES

- Lin, C.C.; Ng, L.T.; Hsu, F.F.; Shieh, D.E.; Chiang, L.C. Cytotoxic effects of *Coptis chinensis* and *Epimedium sagittatum* extracts and their major constituents (berberine, coptisine and icariin) on hepatoma and leukaemia cell growth. *Clin. Exp. Pharmacol. Physiol.*, **2004**, *31*(1-2), 65-69. <http://dx.doi.org/10.1111/j.1440-1681.2004.03951.x> PMID: 14756686
- Zheng, Y.; Feng, W.; Wang, Y.J.; Sun, Y.; Shi, G.; Yu, Q. Galectins as potential emerging key targets in different types of leukemia. *Eur. J. Pharmacol.*, **2019**, *844*, 73-78. <http://dx.doi.org/10.1016/j.ejphar.2018.11.019> PMID: 30452910
- Panagal, M.; S R, S.K.; P, S.; M, B.; M, K.; Gopinathe, V.; Sivakumare, P.; Sekar, D. MicroRNA21 and the various types of myeloid leukemia. *Cancer Gene Ther.*, **2018**, *25*(7-8), 161-166. <http://dx.doi.org/10.1038/s41417-018-0025-2> PMID: 29795410
- Aladag, E.; Haznedaroğlu, İ.C. Current perspectives for the treatment of chronic myeloid leukemia. **2019**, *49*(1), 1-10. <http://dx.doi.org/10.3906/sag-1810-81> PMID: 30761815
- Bazargan, A.; Tam, C.S.; Keating, M.J. Predicting survival in chronic lymphocytic leukemia. *Expert Rev. Anticancer Ther.*, **2012**, *12*(3), 393-403. <http://dx.doi.org/10.1586/era.12.2> PMID: 22369330
- Ilhan, G.; Karakus, S.; Andic, N. Risk factors and primary prevention of acute leukemia. *Asian Pac. J. Cancer Prev.*, **2006**, *7*(4), 515-517. PMID: 17250419
- Heck, J.E.; Park, A.S.; Qiu, J.; Cockburn, M.; Ritz, B. Risk of leukemia in relation to exposure to ambient air toxics in pregnancy and early childhood. *Int. J. Hyg. Environ. Health*, **2014**, *217*(6), 662-668. <http://dx.doi.org/10.1016/j.ijheh.2013.12.003> PMID: 24472648
- Bispo, J.A.B.; Pinheiro, P.S.; Kobetz, E.K. Epidemiology and etiology of leukemia and Lymphoma. *Cold Spring Harb. Perspect. Med.*, **2020**, *10*(6), a034819. <http://dx.doi.org/10.1101/cshperspect.a034819> PMID: 31727680
- Rautenberg, C.; Gerding, U.; Haas, R.; Kobbe, G.; Schroeder, T. Relapse of acute myeloid leukemia after allogeneic stem cell transplantation: Prevention, detection, and treatment. *Int. J. Mol. Sci.*, **2019**, *20*(1), 228. <http://dx.doi.org/10.3390/ijms20010228> PMID: 30626126
- croRNome in cancer and other conditions. *Crit. Rev. Oncol. Hematol.*, **2017**, *116*, 147-158. <http://dx.doi.org/10.1016/j.critrevonc.2017.05.008> PMID: 28693796
- Sun, Y.; Xun, K.; Wang, Y.; Chen, X. A systematic review of the anticancer properties of berberine, a natural product from Chinese herbs. *Anticancer Drugs*, **2009**, *20*(9), 757-769. <http://dx.doi.org/10.1097/CAD.0b013e328330d95b> PMID: 19704371
- Zhao, Y.; Wang, Y.; Ma, S. Racial differences in four leukemia subtypes: Comprehensive descriptive epidemiology. *Sci. Rep.*, **2018**, *8*(1), 548. <http://dx.doi.org/10.1038/s41598-017-19081-4> PMID: 29323237
- Chen, L.S.; Balakrishnan, K.; Gandhi, V. Inflammation and survival pathways: Chronic lymphocytic leukemia as a model system. *Biochem. Pharmacol.*, **2010**, *80*(12), 1936-1945. <http://dx.doi.org/10.1016/j.bcp.2010.07.039> PMID: 20696142
- Jabbour, E.; O'Brien, S.; Konopleva, M.; Kantarjian, H. New insights into the pathophysiology and therapy of adult acute lymphoblastic leukemia. *Cancer*, **2015**, *121*(15), 2517-2528. <http://dx.doi.org/10.1002/cncr.29383> PMID: 25891003
- De Kouchkovsky, I.; Abdul-Hay, M. Acute myeloid leukemia: A comprehensive review and 2016 update. *Blood Cancer J.*, **2016**, *6*(7), e441-e441. <http://dx.doi.org/10.1038/bcj.2016.50> PMID: 27367478
- Talati, C.; Sweet, K. Recently approved therapies in acute myeloid leukemia: A complex treatment landscape. *Leuk. Res.*, **2018**, *73*, 58-66. <http://dx.doi.org/10.1016/j.leukres.2018.09.001> PMID: 30223250
- Finn, L.; Dalovisio, A.; Foran, J. Older patients with acute myeloid leukemia: Treatment challenges and future directions. *Ochsner J.*, **2017**, *17*(4), 398-404. PMID: 29230125
- Mendoza-Nunez, V.M.; Garcia-Martinez, B.I. The effect of 600 mg alpha-lipoic acid supplementation on oxidative stress, inflammation, and RAGE in older adults with type 2 diabetes mellitus. **2019**, *2019*, 3276958. <http://dx.doi.org/10.1155/2019/3276958>
- Hole, P.S.; Darley, R.L.; Tonks, A. Do reactive oxygen species play a role in myeloid leukemias? *Blood*, **2011**, *117*(22), 5816-5826. <http://dx.doi.org/10.1182/blood-2011-01-326025> PMID: 21398578
- Rani, V.; Deep, G.; Singh, R.K.; Palle, K.; Yadav, U.C. Oxidative stress and metabolic disorders: Pathogenesis and therapeutic strategies. *Life Sci.*, **2016**, *148*, 183-193. <http://dx.doi.org/10.1016/j.lfs.2016.02.002> PMID: 26851532
- Samimi, A.; Kalantari, H.; Lorestani, M.Z.; Shirzad, R.; Saki, N. Oxidative stress in normal hematopoietic stem cells and leukemia. *APMIS*, **2018**, *126*(4), 284-294. <http://dx.doi.org/10.1111/apm.12822> PMID: 29575200
- Martindale, J.L.; Holbrook, N.J. Cellular response to oxidative stress: Signaling for suicide and survival. *J. Cell. Physiol.*, **2002**, *192*(1), 1-15. <http://dx.doi.org/10.1002/jcp.10119> PMID: 12115731
- Ariffin, H.; Azanan, M.S.; Abd Ghafar, S.S.; Oh, L.; Lau, K.H.; Thirunavakarasu, T.; Sedan, A.; Ibrahim, K.; Chan, A.; Chin, T.F.; Liew, F.F.; Jeyamogan, S.; Rosli, E.S.; Baharudin, R.; Yap, T.Y.; Skinner, R.; Lum, S.H.; Hainaut, P. Young adult survivors of childhood acute lymphoblastic leukemia show evidence of chronic inflammation and cellular aging. *Cancer*, **2017**, *123*(21), 4207-4214. <http://dx.doi.org/10.1002/cncr.30857> PMID: 28654149
- Binder, S.; Luciano, M.; Horejs-Hoeck, J. The cytokine network in acute myeloid leukemia (AML): A focus on pro- and anti-inflammatory mediators. *Cytokine Growth Factor Rev.*, **2018**, *43*, 8-15. <http://dx.doi.org/10.1016/j.cytogfr.2018.08.004> PMID: 30181021
- Hu, S.; Chen, C.W.; Chen, S.T.; Tsui, K.H.; Tang, T.K.; Cheng, H.T.; Hwang, G.S.; Yu, J.W.; Li, Y.C.; Wang, P.S.; Wang, S.W. Inhibitory effect of berberine on interleukin-2 secretion from PHA-treated lymphocytic Jurkat cells. *Int. Immunopharmacol.*, **2019**, *66*, 267-273. <http://dx.doi.org/10.1016/j.intimp.2018.11.020> PMID: 30502647
- Milata, V.; Svedova, A.; Barbierikova, Z.; Holubkova, E.; Cipakova, I.; Cholujova, D.; Jakubikova, J.; Panik, M.; Jantova, S.; Brezova, V.; Cipak, L. Synthesis and anticancer activity of novel 9-O-

- substituted berberine derivatives. *Int. J. Mol. Sci.*, **2019**, *20*(9), E2169.
<http://dx.doi.org/10.3390/ijms20092169> PMID: 31052469
- [27] Liu, D.; Meng, X.; Wu, D.; Qiu, Z.; Luo, H. A natural isoquinoline alkaloid with antitumor activity: Studies of the biological activities of berberine. *Front. Pharmacol.*, **2019**, *10*, 9.
<http://dx.doi.org/10.3389/fphar.2019.00009> PMID: 30837865
- [28] Cicero, A.F.; Baggioni, A. Berberine and its role in chronic disease. *Adv. Exp. Med. Biol.*, **2016**, *928*, 27-45.
http://dx.doi.org/10.1007/978-3-319-41334-1_2 PMID: 27671811
- [29] Iizuka, N.; Miyamoto, K.; Okita, K.; Tangoku, A.; Hayashi, H.; Yosino, S.; Abe, T.; Morioka, T.; Hazama, S.; Oka, M. Inhibitory effect of Coptidis Rhizoma and berberine on the proliferation of human esophageal cancer cell lines. *Cancer Lett.*, **2000**, *148*(1), 19-25.
[http://dx.doi.org/10.1016/S0304-3835\(99\)00264-5](http://dx.doi.org/10.1016/S0304-3835(99)00264-5) PMID: 10680588
- [30] Kong, W.; Wei, J.; Abidi, P.; Lin, M.; Inaba, S.; Li, C.; Wang, Y.; Wang, Z.; Si, S.; Pan, H.; Wang, S.; Wu, J.; Wang, Y.; Li, Z.; Liu, J.; Jiang, J.D. Berberine is a novel cholesterol-lowering drug working through a unique mechanism distinct from statins. *Nat. Med.*, **2004**, *10*(12), 1344-1351.
<http://dx.doi.org/10.1038/nm1135> PMID: 15531889
- [31] Xue, M.; Yang, M.X.; Zhang, W.; Li, X.M.; Gao, D.H.; Ou, Z.M.; Li, Z.P.; Liu, S.H.; Li, X.J.; Yang, S.Y. Characterization, pharmacokinetics, and hypoglycemic effect of berberine loaded solid lipid nanoparticles. *Int. J. Nanomedicine*, **2013**, *8*, 4677-4687.
<http://dx.doi.org/10.2147/IJN.S51262> PMID: 24353417
- [32] Mirhadi, E.; Rezaee, M.; Malaek-Nikouei, B. Nano strategies for berberine delivery, a natural alkaloid of Berberis. *Biomed. Pharmacother.*, **2018**, *104*, 465-473.
<http://dx.doi.org/10.1016/j.biopha.2018.05.067> PMID: 29793179
- [33] Kumar, A.; Ekavali, ; Chopra, K.; Mukherjee, M.; Pottabathini, R.; Dhull, D.K. Current knowledge and pharmacological profile of berberine: An update. *Eur. J. Pharmacol.*, **2015**, *761*, 288-297.
<http://dx.doi.org/10.1016/j.ejphar.2015.05.068> PMID: 26092760
- [34] Hu, X.; Zhang, Y.; Xue, Y.; Zhang, Z.; Wang, J. Berberine is a potential therapeutic agent for metabolic syndrome via brown adipose tissue activation and metabolism regulation. *Am. J. Transl. Res.*, **2018**, *10*(11), 3322-3329.
 PMID: 30662589
- [35] Feng, X.; Sureda, A.; Jafari, S.; Memariani, Z.; Tewari, D.; Annunziata, G.; Barrea, L.; Hassan, S.T.S.; Šmejkal, K.; Malanik, M.; Sychrová, A.; Barrea, D.; Zibera, L.; Mahomoodally, M.F.; Zengin, G.; Xu, S.; Nabavi, S.M.; Shen, A.Z. Berberine in cardiovascular and metabolic diseases: From mechanisms to therapeutics. *Theranostics*, **2019**, *9*(7), 1923-1951.
<http://dx.doi.org/10.7150/thno.30787> PMID: 31037148
- [36] Imenshahidi, M.; Hosseinzadeh, H. Berberine and barberry (*Berberis vulgaris*): A clinical review. *Phytother. Res.*, **2019**, *33*(3), 504-523.
<http://dx.doi.org/10.1002/ptr.6252> PMID: 30637820
- [37] Zou, K.; Li, Z.; Zhang, Y.; Zhang, H.Y.; Li, B.; Zhu, W.L.; Shi, J.Y.; Jia, Q.; Li, Y.M. Advances in the study of berberine and its derivatives: A focus on anti-inflammatory and anti-tumor effects in the digestive system. *Acta Pharmacol. Sin.*, **2017**, *38*(2), 157-167.
<http://dx.doi.org/10.1038/aps.2016.125> PMID: 27917872
- [38] Li, N.; Gu, L.; Qu, L.; Gong, J.; Li, Q.; Zhu, W.; Li, J. Berberine attenuates pro-inflammatory cytokine-induced tight junction disruption in an *in vitro* model of intestinal epithelial cells. *Eur. J. Pharm. Sci.*, **2010**, *40*(1), 1-8.
<http://dx.doi.org/10.1016/j.ejps.2010.02.001> PMID: 20149867
- [39] Gu, L.; Li, N.; Gong, J.; Li, Q.; Zhu, W.; Li, J. Berberine ameliorates intestinal epithelial tight-junction damage and down-regulates myosin light chain kinase pathways in a mouse model of endotoxemia. *J. Infect. Dis.*, **2011**, *203*(11), 1602-1612.
<http://dx.doi.org/10.1093/infdis/jir147> PMID: 21592990
- [40] Tew, X.N.; Xin Lau, N.J.; Chellappan, D.K.; Madheswaran, T.; Zeeshan, F.; Tambuwala, M.M.; Aljabali, A.A.; Balusamy, S.R.; Perumalsamy, H.; Gupta, G.; Oliver, B.G.; Hsu, A.; Wark, P.; Reddy, K.; Wadhwa, R.; Hansbro, P.M.; Dua, K. Immunological axis of berberine in managing inflammation underlying chronic respiratory inflammatory diseases. *Chem. Biol. Interact.*, **2020**, *317*, 108947.
<http://dx.doi.org/10.1016/j.cbi.2020.108947> PMID: 31968208
- [41] Lu, Z.; He, B.; Chen, Z.; Yan, M.; Wu, L. Anti-inflammatory activity of berberine in non-alcoholic fatty liver disease via the Angptl2 pathway. *BMC Immunol.*, **2020**, *21*(1), 28.
<http://dx.doi.org/10.1186/s12865-020-00358-9> PMID: 32429849
- [42] Eissa, L.A.; Kenawy, H.I.; El-Karef, A.; Elsherbiny, N.M.; El-Mihi, K.A. Antioxidant and anti-inflammatory activities of berberine attenuate hepatic fibrosis induced by thioacetamide injection in rats. *Chem. Biol. Interact.*, **2018**, *294*, 91-100.
<http://dx.doi.org/10.1016/j.cbi.2018.08.016> PMID: 30138605
- [43] Li, Z.; Jiang, T.; Lu, Q.; Xu, K.; He, J.; Xie, L.; Chen, Z.; Zheng, Z.; Ye, L.; Xu, K.; Zhang, H.; Hu, A. Berberine attenuated the cytotoxicity induced by t-BHP via inhibiting oxidative stress and mitochondria dysfunction in PC-12 cells. *Cell. Mol. Neurobiol.*, **2020**, *40*(4), 587-602.
<http://dx.doi.org/10.1007/s10571-019-00756-7> PMID: 31828466
- [44] Dkhil, M.A.; Metwaly, M.S.; Al-Quraishy, S. Berberine improves the intestinal antioxidant status of laboratory mice, *Mus musculus*. *Saudi J. Biol. Sci.*, **2017**, *24*(7), 1567-1573.
<http://dx.doi.org/10.1016/j.sjbs.2015.10.012> PMID: 30294226
- [45] Zhuang, W.; Li, T.; Wang, C.; Shi, X.; Li, Y.; Zhang, S.; Zhao, Z.; Dong, H.; Qiao, Y. Berberine exerts antioxidant effects via protection of spiral ganglion cells against cytomegalovirus-induced apoptosis. *Free Radic. Biol. Med.*, **2018**, *121*, 127-135.
<http://dx.doi.org/10.1016/j.freeradbiomed.2018.04.575> PMID: 29715550
- [46] Zhang, C.; Sheng, J.; Li, G.; Zhao, L.; Wang, Y.; Yang, W.; Yao, X.; Sun, L.; Zhang, Z.; Cui, R. Effects of berberine and its derivatives on cancer: A systems pharmacology review. *Front. Pharmacol.*, **2020**, *10*, 1461-1461.
<http://dx.doi.org/10.3389/fphar.2019.01461> PMID: 32009943
- [47] Muller, P.A.; Vousden, K.H. p53 mutations in cancer. *Nat. Cell Biol.*, **2013**, *15*(1), 2-8.
<http://dx.doi.org/10.1038/ncb2641> PMID: 23263379
- [48] Bieganski, K.T.; Mello, S.S.; Attardi, L.D. Unravelling mechanisms of p53-mediated tumour suppression. *Nat. Rev. Cancer*, **2014**, *14*(5), 359-370.
<http://dx.doi.org/10.1038/nrc3711> PMID: 24739573
- [49] Prokocimer, M.; Molchadsky, A.; Rotter, V. Dysfunctional diversity of p53 proteins in adult acute myeloid leukemia: Projections on diagnostic workup and therapy. *Blood*, **2017**, *130*(6), 699-712.
<http://dx.doi.org/10.1182/blood-2017-02-763086> PMID: 28607134
- [50] Eischen, C.M.; Lozano, G. p53 and MDM2: Antagonists or partners in crime? *Cancer Cell*, **2009**, *15*(3), 161-162.
<http://dx.doi.org/10.1016/j.ccr.2009.02.004> PMID: 19249672
- [51] Van Maerken, T.; Rihani, A.; Van Goethem, A.; De Paepe, A.; Speleman, F.; Vandesompele, J. Pharmacologic activation of wild-type p53 by nutlin therapy in childhood cancer. *Cancer Lett.*, **2014**, *344*(2), 157-165.
<http://dx.doi.org/10.1016/j.canlet.2013.11.002> PMID: 24262662
- [52] Trino, S.; De Luca, L.; Laurenzana, I.; Caivano, A.; Del Vecchio, L.; Martinelli, G.; Musto, P. P53-MDM2 pathway: Evidences for a new targeted therapeutic approach in B-acute lymphoblastic leukemia. *Front. Pharmacol.*, **2016**, *7*, 491.
<http://dx.doi.org/10.3389/fphar.2016.00491> PMID: 28018226
- [53] Liu, J.; Zhu, Z.; Liu, Y.; Wei, L.; Li, B.; Mao, F.; Zhang, J.; Wang, Y.; Liu, Y. MDM2 inhibition-mediated autophagy contributes to the pro-apoptotic effect of berberine in p53-null leukemic cells. *Life Sci.*, **2020**, *242*, 117228.
<http://dx.doi.org/10.1016/j.lfs.2019.117228> PMID: 31881227
- [54] Liu, J.; Zhang, X.; Liu, A.; Liu, S.; Zhang, L.; Wu, B.; Hu, Q. Berberine induces apoptosis in p53-null leukemia cells by down-regulating XIAP at the post-transcriptional level. *Cell. Physiol. Biochem.*, **2013**, *32*(5), 1213-1224.
<http://dx.doi.org/10.1159/000354520> PMID: 24335171
- [55] Zhang, X.; Gu, L.; Li, J.; Shah, N.; He, J.; Yang, L.; Hu, Q.; Zhou, M. Degradation of MDM2 by the interaction between berberine and DAXX leads to potent apoptosis in MDM2-overexpressing cancer cells. *Cancer Res.*, **2010**, *70*(23), 9895-9904.
<http://dx.doi.org/10.1158/0008-5472.CAN-10-1546> PMID: 20935220
- [56] Kapelko-Słowik, K.; Urbaniak-Kujda, D.; Wołowicz, D.; Jazwiec, B.; Dybko, J.; Jakubaszko, J.; Słowik, M.; Kuliczowski, K. Expression of PIM-2 and NF-κB genes is increased in patients with acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL) and is associated with complete remission rate and overall survival. *Postepy Hig. Med. Dosw.*, **2013**, *67*, 553-559.

- <http://dx.doi.org/10.5604/17322693.1052449> PMID: 23752607
- [57] Fox, C.J.; Hammerman, P.S.; Cinalli, R.M.; Master, S.R.; Chodosh, L.A.; Thompson, C.B. The serine/threonine kinase Pim-2 is a transcriptionally regulated apoptotic inhibitor. *Genes Dev.*, **2003**, *17*(15), 1841-1854.
- [58] <http://dx.doi.org/10.1101/gad.1105003> PMID: 12869584
Liu, Z.; Liu, H.; Yuan, X.; Wang, Y.; Li, L.; Wang, G.; Song, J.; Shao, Z.; Fu, R. Downregulation of Pim-2 induces cell cycle arrest in the G₀/G₁ phase via the p53-non-dependent p21 signaling pathway. *Oncol. Lett.*, **2018**, *15*(4), 4079-4086.
- [59] <http://dx.doi.org/10.3892/ol.2018.7865> PMID: 29541172
Wang, Y.; Liu, Y.; Du, X.; Ma, H.; Yao, J. The anti-cancer mechanisms of berberine: A review. *Cancer Manag. Res.*, **2020**, *12*, 695-702.
- [60] <http://dx.doi.org/10.2147/CMAR.S242329> PMID: 32099466
Stagno, F.; Stella, S.; Spitaleri, A.; Pennisi, M.S.; Di Raimondo, F.; Vigneri, P. Imatinib mesylate in chronic myeloid leukemia: Front-line treatment and long-term outcomes. *Expert Rev. Anticancer Ther.*, **2016**, *16*(3), 273-278.
- [61] <http://dx.doi.org/10.1586/14737140.2016.1151356> PMID: 26852913
Valent, P. Imatinib-resistant chronic myeloid leukemia (CML): Current concepts on pathogenesis and new emerging pharmacologic approaches. *Biologics*, **2007**, *1*(4), 433-448.
- [62] PMID: 19707313
Yin, Z.; Huang, G.; Gu, C.; Liu, Y.; Yang, J.; Fei, J. Discovery of berberine that targetedly induce autophagic degradation of both BCR-ABL and BCR-ABL T315I through recruiting LRSAM1 for overcoming imatinib-resistance. *Clin. Cancer Res.*, **2020**, *26*(15), 4040-4053.
- [63] <http://dx.doi.org/10.1158/1078-0432.CCR-19-2460>
Ly, M.; Rentas, S.; Vujovic, A.; Wong, N.; Moreira, S.; Xu, J.; Holzapfel, N.; Bhatia, S.; Tran, D.; Minden, M.D.; Draper, J.S.; Hope, K.J. Diminished AHR signaling drives human acute myeloid leukemia stem cell maintenance. *Cancer Res.*, **2019**, *79*(22), 5799-5811.
- [64] <http://dx.doi.org/10.1158/0008-5472.CAN-19-0274> PMID: 31519687
Gentil, M.; Hugues, P.; Desterke, C.; Telliam, G.; Sloma, I.; Souza, L.E.B.; Baykal, S.; Artus, J.; Griscelli, F.; Guerci, A.; Johnson-Ansah, H.; Foudi, A.; Bennaceur-Griscelli, A.; Turhan, A.G. Aryl hydrocarbon receptor (AHR) is a novel druggable pathway controlling malignant progenitor proliferation in chronic myeloid leukemia (CML). *PLoS One*, **2018**, *13*(8), e0200923.
- [65] <http://dx.doi.org/10.1371/journal.pone.0200923> PMID: 30091999
Mohammadi, S.; Seyedhoseini, F.S.; Asadi, J.; Yazdani, Y. Effects of berberine on the secretion of cytokines and expression of genes involved in cell cycle regulation in THP-1 monocytic cell line. *Iran. J. Basic Med. Sci.*, **2017**, *20*(5), 530-537.
- [66] PMID: 28656088
Wang, L.; Chen, B.; Lin, M.; Cao, Y.; Chen, Y.; Chen, X.; Liu, T.; Hu, J. Decreased expression of nucleophosmin/B23 increases drug sensitivity of adriamycin-resistant Molt-4 leukemia cells through mdr-1 regulation and Akt/mTOR signaling. *Immunobiology*, **2015**, *220*(3), 331-340.
- [67] <http://dx.doi.org/10.1016/j.imbio.2014.10.015> PMID: 25457413
Zhang, M.J.; Ding, Y.L.; Xu, C.W.; Yang, Y.; Lian, W.X.; Zhan, Y.Q.; Li, W.; Xu, W.X.; Yu, M.; Ge, C.H.; Ning, H.M.; Li, C.Y.; Yang, X.M. Erythroid differentiation-associated gene interacts with NPM1 (nucleophosmin/B23) and increases its protein stability, resisting cell apoptosis. *FEBS J.*, **2012**, *279*(16), 2848-2862.
- [68] <http://dx.doi.org/10.1111/j.1742-4658.2012.08663.x> PMID: 22712502
Wu, H.L.; Hsu, C.Y.; Liu, W.H.; Yung, B.Y. Berberine-induced apoptosis of human leukemia HL-60 cells is associated with down-regulation of nucleophosmin/B23 and telomerase activity. *Int. J. Cancer*, **1999**, *81*(6), 923-929.
- [69] [http://dx.doi.org/10.1002/\(SICI\)1097-0215\(19990611\)81:6<923::AID-IJC14>3.0.CO;2-D](http://dx.doi.org/10.1002/(SICI)1097-0215(19990611)81:6<923::AID-IJC14>3.0.CO;2-D) PMID: 10362140
Fu, S.; Ni, S.; Wang, D.; Fu, M.; Hong, T. Berberine suppresses mast cell-mediated allergic responses via regulating FcεRI-mediated and MAPK signaling. *Int. Immunopharmacol.*, **2019**, *71*, 1-6.
- [70] <http://dx.doi.org/10.1016/j.intimp.2019.02.041> PMID: 30861392
Lin, C.C.; Lin, S.Y.; Chung, J.G.; Lin, J.P.; Chen, G.W.; Kao, S.T. Down-regulation of cyclin B1 and up-regulation of wee1 by berberine promotes entry of leukemia cells into the G₂/M-phase of the cell cycle. *Anticancer Res.*, **2006**, *26*(2A), 1097-1104.
- [71] PMID: 16619512
He, Z.W.; Zhao, X.Y.; Xu, R.Z.; Wu, D. Effects of berbamine on growth of leukemia cell line NB4 and its mechanism. *Zhejiang Da Xue Xue Bao Yi Xue Ban*, **2006**, *35*(2), 209-214.
- [72] PMID: 16610091
Diab, S.; Fidanzi, C.; Léger, D.Y.; Ghezali, L.; Millot, M.; Martin, F.; Azar, R.; Esseily, F.; Saab, A.; Sol, V.; Diab-Assaf, M.; Liagre, B. *Berberis libanotica* extract targets NF-κB/COX-2, PI3K/Akt and mitochondrial/caspase signalling to induce human erythroleukemia cell apoptosis. *Int. J. Oncol.*, **2015**, *47*(1), 220-230.
- [73] <http://dx.doi.org/10.3892/ijo.2015.3012> PMID: 25997834
Okubo, S.; Uto, T.; Goto, A.; Tanaka, H.; Nishioku, T.; Yamada, K.; Shoyama, Y. Berberine induces apoptotic cell death via activation of caspase-3 and -8 in HL-60 human leukemia cells: Nuclear localization and structure-activity relationships. *Am. J. Chin. Med.*, **2017**, *45*(7), 1497-1511.
- [74] <http://dx.doi.org/10.1142/S0192415X17500811> PMID: 29025293
Li, H.; Guo, L.; Jie, S.; Liu, W.; Zhu, J.; Du, W.; Fan, L.; Wang, X.; Fu, B.; Huang, S. Berberine inhibits SDF-1-induced AML cells and leukemic stem cells migration via regulation of SDF-1 level in bone marrow stromal cells. *Biomed. Pharmacother.*, **2008**, *62*(9), 573-578.
- [75] <http://dx.doi.org/10.1016/j.biopha.2008.08.003> PMID: 18805669
Lin, C.C.; Kao, S.T.; Chen, G.W.; Ho, H.C.; Chung, J.G. Apoptosis of human leukemia HL-60 cells and murine leukemia WEHI-3 cells induced by berberine through the activation of caspase-3. *Anticancer Res.*, **2006**, *26*(1A), 227-242.
- [76] PMID: 16475703
Chung, J.G.; Chen, G.W.; Hung, C.F.; Lee, J.H.; Ho, C.C.; Ho, H.C.; Chang, H.L.; Lin, W.C.; Lin, J.G. Effects of berberine on arylamine N-acetyltransferase activity and 2-aminofluorene-DNA adduct formation in human leukemia cells. *Am. J. Chin. Med.*, **2000**, *28*(2), 227-238.
- [77] <http://dx.doi.org/10.1142/S0192415X00000271> PMID: 10999441
Lin, S.S.; Chung, J.G.; Lin, J.P.; Chuang, J.Y.; Chang, W.C.; Wu, J.Y.; Tyan, Y.S. Berberine inhibits arylamine N-acetyltransferase activity and gene expression in mouse leukemia L 1210 cells. *Phytomedicine*, **2005**, *12*(5), 351-358.
- [78] <http://dx.doi.org/10.1016/j.phymed.2003.11.008> PMID: 15957369
Lin, C.C.; Kao, S.T.; Chen, G.W.; Chung, J.G. Berberine decreased N-acetylation of 2-aminofluorene through inhibition of N-acetyltransferase gene expression in human leukemia HL-60 cells. *Anticancer Res.*, **2005**, *25*(6B), 4149-4155.
- [79] PMID: 16309210
Lu, Y.-C.; Lin, Q.; Luo, G.-S.; Dai, Y.-Y. Solubility of berberine chloride in various solvents. *J. Chem. Eng. Data*, **2006**, *51*(2), 642-644.
- [80] <http://dx.doi.org/10.1021/je0504360>
Ge, L.; Bao, A.; Yang, K. Determination and correlation of the solubility for berberine chloride in pure imidazolium-based ionic liquids. *J. Chem. Eng. Data*, **2016**, *61*(5), 1829-1835.
- [81] <http://dx.doi.org/10.1021/acs.jced.5b01017>
Zhu, X.; Sun, Y.; Zhang, C.; Liu, H. Effects of berberine on a rat model of chronic stress and depression via gastrointestinal tract pathology and gastrointestinal flora profile assays. *Mol. Med. Rep.*, **2017**, *15*(5), 3161-3171.
- [82] <http://dx.doi.org/10.3892/mmr.2017.6353> PMID: 28339024
Xiao, D.; Liu, Z.; Zhang, S.; Zhou, M.; He, F.; Zou, M.; Peng, J.; Xie, X.; Liu, Y.; Peng, D. Berberine derivatives with different pharmacological activities via structural modifications. *Mini Rev. Med. Chem.*, **2018**, *18*(17), 1424-1441.
- [83] <http://dx.doi.org/10.2174/1389557517666170321103139> PMID: 28325147
Sagar, B. *In-vitro* production of berberine in cultures of *Berberis aristata* and pharmacological investigation for anti-hepatotoxic activity. *J. Pharm. Pharmacol.*, **2006**, *58*.
- [84] Guaman Ortiz, L.M.; Tillhon, M.; Parks, M.; Dutto, I.; Prosperi, E.; Savio, M.; Arcamone, A.G.; Buzzetti, F.; Lombardi, P.; Scovassi, A.I. Multiple effects of berberine derivatives on colon cancer cells. *BioMed Res. Int.*, **2014**, *2014*, 924585.
- <http://dx.doi.org/10.1155/2014/924585> PMID: 25045712

- [85] Park, K.D.; Lee, J.H.; Kim, S.H.; Kang, T.H.; Moon, J.S.; Kim, S.U. Synthesis of 13-(substituted benzyl) berberine and berberrubine derivatives as antifungal agents. *Bioorg. Med. Chem. Lett.*, **2006**, *16*(15), 3913-3916.
<http://dx.doi.org/10.1016/j.bmcl.2006.05.033> PMID: 16730982
- [86] Huang, L.; Luo, Z.; He, F.; Shi, A.; Qin, F.; Li, X. Berberine derivatives, with substituted amino groups linked at the 9-position, as inhibitors of acetylcholinesterase/butyrylcholinesterase. *Bioorg. Med. Chem. Lett.*, **2010**, *20*(22), 6649-6652.
<http://dx.doi.org/10.1016/j.bmcl.2010.09.013> PMID: 20880702
- [87] Cui, J.S.; Xu, F.; Pang, J.Y.; Chen, W.H.; Jiang, Z.H. Synthesis and DNA-binding affinities of protoberberine-based multivalent agents. *Chem. Biodivers.*, **2010**, *7*(12), 2908-2916.
<http://dx.doi.org/10.1002/cbdv.200900386> PMID: 21162004
- [88] Pang, J.Y.; Long, Y.H.; Chen, W.H.; Jiang, Z.H. Amplification of DNA-binding affinities of protoberberine alkaloids by appended polyamines. *Bioorg. Med. Chem. Lett.*, **2007**, *17*(4), 1018-1021.
<http://dx.doi.org/10.1016/j.bmcl.2006.11.037> PMID: 17127054
- [89] Pang, J.Y.; Qin, Y.; Chen, W.H.; Luo, G.A.; Jiang, Z.H. Synthesis and DNA-binding affinities of monomodified berberines. *Bioorg. Med. Chem.*, **2005**, *13*(20), 5835-5840.
<http://dx.doi.org/10.1016/j.bmc.2005.05.048> PMID: 15993616
- [90] Krishnan, P.; Bastow, K.F. The 9-position in berberine analogs is an important determinant of DNA topoisomerase II inhibition. *Anticancer Drug Des.*, **2000**, *15*(4), 255-264. PMID: 11200501
- [91] Iwasa, K.; Kamiguchi, M.; Ueki, M.; Taniguchi, M. Antibacterial activity and structure-activity relationships of berberine analogs. *Eur. J. Med. Chem.*, **1996**, *31*(6), 469-478.
[http://dx.doi.org/10.1016/0223-5234\(96\)85167-1](http://dx.doi.org/10.1016/0223-5234(96)85167-1)
- [92] Inoue, K.; Kulsum, U.; Chowdhury, S.A.; Fujisawa, S.; Ishihara, M.; Yokoe, I.; Sakagami, H. Tumor-specific cytotoxicity and apoptosis-inducing activity of berberines. *Anticancer Res.*, **2005**, *25*(6B), 4053-4059. PMID: 16309199
- [93] Tang, P.; Cheng, W.; He, X.; Zhang, Q.; Zhong, J.; Lu, X.; Liu, S.; Zhong, L. Raman spectrum spectral imaging revealing the molecular mechanism of berberine-induced Jurkat cell apoptosis and the receptor-mediated berberine delivery system. *Biomed. Opt. Express*, **2019**, *10*(4), 1581-1600.
<http://dx.doi.org/10.1364/BOE.10.001581> PMID: 31061758
- [94] Jin, L.; Liao, H.J.; Zhang, M.Y.; Liu, Q.Y.; Wang, Y.F. Effect of berberine on the differentiation and apoptosis of K562 cell line. *Zhong Yao Cai*, **2009**, *32*(3), 384-388. PMID: 19565717
- [95] Pazhang, Y.; Ahmadian, S.; Mahmoudian, M.; Shafieezadeh, M. Berberine-induced apoptosis via decreasing the survivin protein in K562 cell line. *Med. Oncol.*, **2011**, *28*(4), 1577-1583.
<http://dx.doi.org/10.1007/s12032-010-9586-0> PMID: 20517657
- [96] Ma, Y.; Ou, T.M.; Tan, J.H.; Hou, J.Q.; Huang, S.L.; Gu, L.Q.; Huang, Z.S. Quinolono-benzo-[5, 6]-dihydroisoquinolium compounds derived from berberine: A new class of highly selective ligands for G-quadruplex DNA in c-myc oncogene. *Eur. J. Med. Chem.*, **2011**, *46*(5), 1906-1913.
<http://dx.doi.org/10.1016/j.ejmech.2011.02.020> PMID: 21392861
- [97] Jantová, S.; Cipák, L.; Cernáková, M.; Kost'álová, D. Effect of berberine on proliferation, cell cycle and apoptosis in HeLa and L1210 cells. *J. Pharm. Pharmacol.*, **2003**, *55*(8), 1143-1149.
<http://dx.doi.org/10.1211/002235703322277186> PMID: 12956905
- [98] Yu, F.S.; Yang, J.S.; Lin, H.J.; Yu, C.S.; Tan, T.W.; Lin, Y.T.; Lin, C.C.; Lu, H.F.; Chung, J.G. Berberine inhibits WEHI-3 leukemia cells *in vivo*. *In Vivo*, **2007**, *21*(2), 407-412. PMID: 17436595
- [99] Kuo, C.L.; Chou, C.C.; Yung, B.Y. Berberine complexes with DNA in the berberine-induced apoptosis in human leukemic HL-60 cells. *Cancer Lett.*, **1995**, *93*(2), 193-200.
[http://dx.doi.org/10.1016/0304-3835\(95\)03809-B](http://dx.doi.org/10.1016/0304-3835(95)03809-B) PMID: 7621428
- [100] Kettmann, V.; Kosfálová, D.; Jantová, S.; Cernáková, M.; Drimal, J. *In vitro* cytotoxicity of berberine against HeLa and L1210 cancer cell lines. *Pharmazie*, **2004**, *59*(7), 548-551. PMID: 15296093
- [101] Liu, J.; Liu, P.; Xu, T.; Chen, Z.; Kong, H.; Chu, W.; Wang, Y.; Liu, Y. Berberine induces autophagic cell death in acute lymphoblastic leukemia by inactivating AKT/mTORC1 signaling. *Drug Des. Devel. Ther.*, **2020**, *14*, 1813-1823.
<http://dx.doi.org/10.2147/DDDT.S239247> PMID: 32494123
- [102] Yin, Z.; Huang, G.; Gu, C.; Liu, Y.; Yang, J.; Fei, J. Discovery of berberine that targetedly induces autophagic degradation of both BCR-ABL and BCR-ABL T3151 through recruiting LRSAM1 for overcoming imatinib resistance. *Clin. Cancer Res.*, **2020**, *26*(15), 4040-4053.
<http://dx.doi.org/10.1158/1078-0432.CCR-19-2460> PMID: 32098768